



Transporting Children

Driver: _____

Bus # _____

Date: _____

Child's Name	Signs of COVID-19 N or Y	Morning Run			Evening Run	
		TEMP	Time On	Time Off	Time On	Time Off

- **COVID-19 Screening Questions:**
 - ✓ A new fever (100.4°F or higher) or a sense of having a fever?
 - ✓ Any new cough that cannot be attributed to another health condition?
 - ✓ New shortness of breath that cannot be attributed to another health condition?
 - ✓ New chills that cannot be attributed to another health condition?
 - ✓ Any new sore throat that cannot be attributed to another health condition?
 - ✓ New muscle aches (myalgia) that cannot be attributed to another health condition or specific activity (such as physical exercise)?
 - ✓ Have been in close contact or cared for someone with COVID 19?


Bus Driver Signature

Classroom Staff/Teacher

VDH Algorithm for Evaluating a Child with COVID-19 Symptoms or Exposure (August 24, 2020)

For Parents and Guardians


***Symptoms** of COVID-19 include fever (≥100.4°F) or chills, fatigue (more tired than usual), headache, muscle aches, cough, nasal congestion or runny nose, new loss of taste or smell, sore throat, shortness of breath or difficulty breathing, abdominal pain, diarrhea, nausea or vomiting, new onset poor appetite or poor feeding.



For Schools and Childcare Facilities

****Close contact** means being within 6 feet of a person with COVID-19 for 15 minutes or more or direct exposure to respiratory secretions


***** ≥10 cases per 100,000 population** Currently all of Virginia. This will be updated with a link to data on local transmission when available.



For Healthcare Providers

Clinical Evaluation for Children with Symptoms of COVID-19* (www.cdc.gov/coronavirus/2019-ncov/hcp/pediatric-hcp.html)

† Testing – PCR or antigen (Ag) testing is acceptable. If an Ag detection test is negative and there is a high clinical suspicion of COVID-19, confirm with PCR, ideally within 2 days of the initial Ag test. If RT-PCR testing is not available, clinical discretion can be used in whether to recommend the patient isolate.



Return to School and Childcare

